

West Nile Virus Seroprevalence in Manisa Province: A Population-based Study

Abdurrahman Gülmez¹ , Ahmet Naci Emecen² , Mestan Emek³ , Belgin Ünal² , Koray Ergünay⁴ , İbrahim Mehmet Ali Öktem⁵ , Özgen Alpay Özbek⁶ 

¹ Medical Microbiology Laboratory, İstanbul Başakşehir Cam ve Sakura Hospital, İstanbul, Turkey

² Department of Public Health, Dokuz Eylül University School of Medicine, İzmir, Turkey

³ Department of Public Health, Akdeniz University School of Medicine, Antalya, Turkey

⁴ Department of Medical Microbiology, Virology Unit, Hacettepe University School of Medicine, Ankara, Turkey

⁵ Department of Medical Microbiology, Virology Unit, Dokuz Eylül University School of Medicine, İzmir, Turkey

⁶ Department of Medical Microbiology, Dokuz Eylül University School of Medicine, İzmir, Turkey

ABSTRACT

Objective: West Nile Virus (WNV), which causes widespread outbreaks in different parts of the world, is a risk to public health in Turkey, too. Community-based study data are needed to identify measures against possible outbreaks. This study aimed to determine the seroprevalence of community-based WNV in Manisa and to investigate the relationship between sociodemographic and socioeconomic variables.

Methods: We included individuals older than two years of age (N=1,317,917) registered in the Manisa Province Family Medicine Information System. Selected participants (n=1233) were determined by a simple random sampling method. Specific IgG antibodies against WNV were investigated in serum samples using a commercial ELISA test (Euroimmun, Germany). The relationship between age, gender, location, education and income level, occupation, population density, altitude, the location of the toilet in the house, and the presence of hypertension, diabetes mellitus and cardiovascular disease variables were analyzed by chi-square, Fisher's exact test and t-test. Adjusted odds ratio (OR) with 95% confidence interval (CI) for each variable were calculated by the logistic regression method to explain potential risks.

Results: WNV IgG antibodies were detected in 47 (3.8%) sera samples by ELISA. Seroprevalence was significantly correlated with independent variables of advanced age, presence of hypertension, diabetes mellitus and cardiovascular disease, low level of education and income, living in low altitude areas and the location of the toilet. In multivariate analysis; age (every one-year increase) (OR:1.05; 95% CI:1.02-1.07; $p < 0.001$), equivalent annual income per capita below 3265 TL (OR:3.21; 95% CI: 1.53-6.73; $p=0.002$), and living areas below 132 meters altitude (OR=3.21; 95% CI 1.26-8.15; $p=0.014$) were found to be the risk factors for WNV seropositivity.

Conclusion: In Manisa province, WNV IgG seroprevalence was detected as 3.8% with ELISA method. Older age, low income and living in regions with a low altitude were found to be associated with increased seropositivity significantly.

Keywords: West Nile virus, seroprevalence, ELISA

Corresponding Author:
Abdurrahman Gülmez

E-mail:
drgulmez@gmail.com

Received: November 22, 2021

Accepted: March 05, 2022

Published: June 13, 2022

Suggested citation:

Gülmez A, Emecen AN, Emek M, Ünal B, Ergünay K, Öktem İMA, et al. West Nile virus seroprevalence in Manisa province: A population-based study. *Infect Dis Clin Microbiol*. 2022;2:107-15.

DOI: 10.36519/idcm.2022.102



INTRODUCTION

West Nile Virus (WNV) is an enveloped RNA virus with icosahedral symmetry and with a size of 45-50 nm belonging to the family *Flaviviridae*, genus *Flavivirus* (1). The virus, which was first isolated from a woman living in the West Nile region of Uganda in 1937, was noted with consecutive outbreaks in North Africa, Southern Europe, Israel and Russia in the 1990s and its spread has been accelerated with its passage to North America at the beginning of the 2000s. WNV outbreaks have been seen frequently especially in the Middle East and Europe in recent years (2, 3).

Transmission usually occurs with the mediation of the mosquitoes of *Culex* genus that feed on infected birds. Thus, birds are the natural sources and reservoir hosts of WNV. Also, it has been reported that transmission via blood and organ transplantation, transplacental route and breast milk was possible (4, 5).

In Turkey, The highest number of WNV cases were detected in 2010. In the study conducted by Kalaycioglu et al. between 2010 and 2011, 47 cases were diagnosed with WNV, and it was reported that the virus could be endemic in the western regions of Turkey (6).

Turkey is vulnerable to WNV cases and outbreaks due to the presence of infectious vectors and its geographical location on the flight paths of migratory birds.

This community-based study aimed to determine the seroprevalence of WNV in serum samples col-

lected from the whole Manisa province in 2014 and the relation of this seroprevalence with sociodemographic and socioeconomic variables.

MATERIALS AND METHODS

Study Population

The sera and data were collected between March 18, 2014, and June 22, 2014, for "Determination of the seroprevalence of some vaccine-preventable diseases in Manisa, 2014" Research Project (BAP) of Dokuz Eylül University.

This is a cross-sectional seroprevalence study and its population consists of all individuals over the age of two who are registered in the Manisa Province Family Medicine Information System (FMIS) (N=1,317,917). With OpenEpi software, version 3.01 (www.openepi.com), the smallest sample size was determined to be 1337 people with 2% expected prevalence, 0.75% standard deviation and 95% confidence interval. By considering the problems that may be encountered during data collection or problems such as the possibility of being unable to reach the people in the sample, insufficient serum, hemolysis, and lipolysis, 30% backup was added to this number and the sample size was calculated to be 1740. Of the targeted 1740 people, 1260 (72.4%) people accepted to be included in the study and 1255 people gave a blood sample for serum.

The study subjects were selected by using a simple random sampling method with the help of a database (Oracle, Austin, Texas, U.S.). The similarity of the demographic characteristics of the selected people such as age, sex and place of residence with those of the population was evaluated. The people who were chosen as a result of sample selection were invited to the Family Medicine Centers (FMC) in which they were registered by appointment, between March 2014 and June 22, 2014, by the FMC staff. Data collection forms were filled by pollsters by face-to-face interview method. During the interview, 10 ml venous blood sample was obtained by a nurse or midwife. The collected blood samples were sent to Manisa Public Health Laboratory on the same day and were frozen at -20 °C after being centrifuged and aliquoted. The serums were weekly transferred to Dokuz Eylül University Medical Microbiology Laboratory by cold chain and stored at -80 °C.

HIGHLIGHTS

- Seroprevalence of WNV IgG was found to be 3.8% in Manisa Province.
- Advanced age, low income and living in low altitude areas have a significant effect on seropositivity.
- Underlying disease, living in a rural area or population density were not found to be associated with seropositivity.

Questionnaire and Data Collection

The ages of the participants were classified within groups of 10-year intervals, their locations as “periphery” or “central district” in relation to the FMC region, and their educational statuses as “illiterate”, “literate”, “elementary school”, “middle school”, “high school”, “university and above”. Presence and type (hypertension [HT], diabetes mellitus [DM], cardiovascular disease [CVD]) of chronic disease if present were questioned. The location of the toilet (whether it was in or out of the house) was questioned for vector exposure. The monthly income of the participants was asked with an open-ended question and the yearly equivalized income per person pertaining to the year 2014 was calculated to be 3265 TL according to the equivalized income scale developed by Organization for Economic

Co-operation and Development (OECD). People with incomes lower than this income were considered to be low-income and those who had higher incomes than this were considered to be high income.

The mean altitudes of the districts and their areas in terms of square kilometers were obtained from the official website of Republic of Turkey Ministry of National Defence, General Directorate of Mapping. The population data of cities and districts pertaining to 2013 were obtained from the database of TURKSTAT website (Turkish Statistical Institute) (7-8). The provincial median altitude value was calculated to be 132 meters (m), the districts were separated into two as being below or above this value. Population densities were calculated separately by dividing the district populations by their areas and the provincial median was found to be 80 people/km² and the districts were separated into two groups.

Laboratory Analysis of Samples

In the study, the specific antibody responses to WNV in the serum samples of the participants were examined by a commercial ELISA (Enzyme-linked immunosorbent assay) method according to the manufacturer’s instructions (Euroimmun, Lubeck, Germany). The antigen source used in the test is a recombinant, detergent-extracted glycoprotein E of WNV from the membrane fraction of human cells. Quantitative measurements were done for IgG-type antibodies using optical density and

Table 1. WNV IgG antibody results.

WNV IgG	Positive (+)	Borderline (-/+)	Negative	Total
n	47	1	1185	1233
%	3.8	0.1	96.1	100

standard calibrators. All indeterminate results were considered to be negative.

The statistical analysis was performed by using the Statistical Package for Social Sciences (SPSS) 22.0 (IBM Corp., Armonk, NY, USA). Chi-square test was used to detect the association of antibody positivity with gender, age group, presence of chronic disease, hypertension, diabetes, cardiovascular diseases, educational status, altitude, population density, and location of the toilet in the house. By including the significant variables logistic regression was performed for the prediction of seropositivity. Analyses were conducted within 95% confidence interval (CI) and $p < 0.05$ values were set as significance. Dokuz Eylül University Non-Invasive Research Ethics Committee approved the study on August 13, 2018, with the decision number of “2018/17-07”.

RESULTS

Of the 1255 people who consented to be included, 22 were excluded because of an insufficient sample, so the results were analyzed for 1233 people.

In the study group, 591 (47.9%) were males, 642 (52.1) were females and their ages varied between 2 and 82 (mean age 36.72 ± 21.09).

In the study, WNV IgG was positive in 3.8% (47/1233) and indeterminate in 0.1% (1/1233) (Table 1). Because the sera detected to be positive as a result of antibodies were not confirmed with the plaque reduction neutralization test (PRNT) method, the results were interpreted as probable antibody positivity and analysis findings were calculated only over positive sera.

In this study, no association was found between gender and seroprevalence ($p=0.46$), but a significant correlation was found between seroprevalence and age groups ($p < 0.001$). This difference was de-




Figure 1. WNV IgG seroprevalence in age groups (%).




Figure 2. Mean values of WNV IgG antibody level (RU/mL) in age groups ($p=0.6$).

tected among individuals older than 40-49 age group (Figure 1). WNV IgG antibody level (RU/mL) was detected to be similar in all age groups ($p=0.6$) (Figure 2).

WNV IgG seroprevalence was found to be significantly higher among the people with HT, DM and CVD in comparison to those who did not have these diseases (Table 2).

WNV IgG seroprevalence was significantly found to be higher in adults who were illiterate, merely literate or elementary school graduates. This difference was observed to become more prominent as the level of education decreased ($p<0.001$). There was

Table 2. Association between participants' disease histories and antibody results.

	WNV IgG			
	Positive	Negative	Seroprevalence (%)	<i>p</i>
Chronic disease (any)				
+	21	241	8	<0.001
-	26	924	2.7	
Hypertension				
+	18	179	9.1	<0.001
-	29	986	2.9	
Diabetes mellitus				
+	8	88	8.3	0.01
-	38	1076	3.4	
Cardiovascular disease				
+	20	210	8.7	<0.001
-	27	954	3.4	

no difference in seroprevalence between central and peripheral FMC regions. When the examination was done according to annual adjusted income per person; WNV IgG seroprevalence was found to be significantly higher for those with low income ($p=0.03$) (Table 3).

The WNV IgG seroprevalence was found to be significantly higher among the participants who lived in the houses where the toilets were located outside of the house or toilets both inside and outside the house than those with toilets inside the house ($p=0.01$).

The people in the study group resided in the 17 districts within the Manisa province borders. The WNV IgG seroprevalence was detected to be higher in some districts. These districts were marked on geographical maps and it was seen that they were located in Gediz and Bakırçay river beds with relatively lower altitudes (Figure 3).

The WNV IgG seroprevalence in the districts with a median altitude value below 132 m was found to be significantly higher than those with a median altitude of more than 132 m ($p=0.02$). No difference

Table 3. Factors affecting WNV IgG seroprevalence (univariate-multivariate regression analysis).

Independent variable	Univariate analysis			Multivariate analysis		
	OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>
Age						
Continuous	1.05	1.04-1.07	<0.001	1.05	1.03-1.07	<0.001
Sex						
Female/Male	1.25	0.7-2.23	0.463			
Living place						
Periphery/ Central	1.54	0.81-2.92	0.185			
Educational status						
≥Middle school / ≤ Elementary school	3.8	1.69-8.57	0.001	1.38	0.54-3.54	0.5
Annual equivalized income (TL)						
>3265 / ≤ 3265	2	1.05-3.77	0.048	3.21	1.53-6.73	0.002
Location of the toilet						
Inside/ Outside	2.21	1.21-4.04	0.013	1.59	0.80-3.17	0.18
Chronic diseases						
Hypertension						
-/+	3.09	1.71-5.6	<0.001			
-/+	3.42	1.86-6.29	<0.001	0.99	0.19-5.14	0.995
Diabetes mellitus						
-/+	2.57	1.17-5.69	0.023	1.48	0.59-3.70	0.4
Cardiovascular disease						
-/+	3.37	1.85-6.16	<0.001	1.01	0.21-4.96	0.986
Altitude (m)						
>132 / ≤132	2.32	1.07-5.01	0.027	3.21	1.27-8.16	0.014
Population density (person/km²)						
≤80 / >80	1.01	0.56-1.81	0.878			

OR: Odds Ratio, CI: Confidence Interval,

was detected in seroprevalence between districts with a population density higher than 80 people/km² and those with a population density less than 80 people/km² (*p*=0.87).

In the multivariate analysis, one unit increase in age caused a significant increase of 1.05 times, an annual adjusted income per person less than 3265 TL increased it by 3.21 times and an altitude of 132 m or below increased it by 3.21 times (Table 3).

DISCUSSION

It has been demonstrated in the studies conducted under experimental conditions that in recent years not only the prevalence but also the virulence has increased (9-12). However, difficulties caused by cross-reactions in the laboratory diagnosis of the disease and limitations in treatment options continue and there is no approved vaccine that can be used in humans (13). Thus, there is a need for epidemiologic data that are to be obtained from the




Figure 3. Manisa province topographic map and WNV IgG seroprevalence map in its districts (%).

studies in the world and our country and for protective measures that will be created with the use of this data. This study is a community-based WNV seroprevalence study and epidemiological data for Manisa province was obtained.

West Nile virus seroprevalence was reported as 11.0% in the 15-65 age interval in Iran by using the cluster sampling method and ELISA (14), it was reported as 24% in Egypt by using a random sampling method and PRNT (15). In a study conducted in Greece using a random sampling method from the general population and ELISA-PRNT tests, the mean seroprevalence in all age groups was reported as 2.1% and 1.5% respectively (16). In the present study, the surveillance was found to be 3.8% for Manisa province.

In three different studies conducted with healthy blood donors in our country between 2010 and 2015 with ELISA method, WNV seropositivity in 2821, 2516, 1200 serum samples were found as 2.4%, 0.99% and 1.6% respectively. While no virus was detected in the first of these studies with polymerase chain reaction (PCR), specific antibody presence was confirmed in nearly half of the samples that were detected to be positive with PRNT method in the other two studies (17-19). Blood donors usually consist of healthy young adults. However, in case a disease or infection finding is detected in the pre-donation examination of the blood donor candidates, these people are excluded from donation permanently or temporarily before blood is taken

(20). WNV seroprevalence was detected to be low in blood donors in this study. This might be explained by the presence of people who cannot donate blood due to medical reasons and people older than 65 years of.

According to the annual data of the European Center for Disease Prevention and Control (ECDC), WNV infection is seen more frequently in men when compared to women (M/W ratio:1.8) (21). In the present study, while WNV IgG seroprevalence is similarly higher for men (M/W ratio:1.23), no significant difference was detected between the sexes ($p=0.46$).

In previous studies, it was reported that being over 50 years of age was a risk factor for being seropositive, that it increased seropositivity by 3.5 to 5.2 times and that seroprevalence also increased by increasing age (22-23). Similarly, we detected an association between increased age and increased WNV IgG seroprevalence significantly ($p<0.001$).

The results revealing a higher WNV seroprevalence in the study region may be due to the geographical characteristics of Manisa province. A big part of the province is located in the Gediz river basin and three dams and many streams as well as two important rivers, namely Gediz and Bakircay, are located in this river basin (24). In this region, the genera of mosquitoes and their seasonal change throughout Manisa province, 3 of the *Culex* genus (*Culex pipens*, *Culex martinii*, *Culex deserticola*) were

detected to be prevalent and *C. pipens* and *C. martinii* were reported as dominant species (25). Also, about 211 different bird species have been observed in this basin and changes are seen in their numbers during the migration seasons (26).

The presence of vector mosquito species is influenced by geographical conditions and altitude. In regions with low altitudes, vectors are more prevalent when compared with those in high altitudes (27). In previous studies, it was observed that viral spread was more frequently seen in regions with low altitudes (28) and that the risk decreased on mountainside (29). When the districts in this study were separated into two groups as those with a median altitude value below 132 m and those with a median altitude value over 132 m, WNV IgG seroprevalence was observed to be significantly higher in the districts with an altitude below 132 m ($p=0.02$). This significance continued also in the multivariate analysis and seropositivity risk was detected to be 3.2 times higher in people living in regions with an altitude below 132 m.

It has been demonstrated that increases in population density and housing population caused vector spread and increase (30,31) and that this situation increased and influenced the geographical distribution of the virus (29). The districts in the study region were separated into two groups as those with a population density over 80 people/km² value and below 80 people/km² value but no difference was detected in seroprevalence in terms of population density ($p=0.97$).

In low-income regions, increased vector activity (30), WNV seroprevalence (32) and neuroinvasive disease development risk (33) were observed to be higher than those in other regions. It has been reported that the presence of old houses in these regions could also be supplying appropriate reproduction areas for vectors (30). In the study, adjusted income per person was calculated with the OECD formula for 2014; WNV IgG seroprevalence was found to be significantly higher for people with an income below 3265 TL ($p=0.03$). The location of the toilet in the house was considered as constituting a risk factor for vector theme in addition to poverty indicator; WNV IgG seroprevalence was found to

be higher for people with toilets outside the house in comparison to people with toilets in the house ($p=0.01$) (Table 3). However, in the multivariate model, it was observed that this significance disappeared. In addition, it was observed that the central and periphery of the living area did not affect the seroprevalence. When these data in the study were evaluated together, they suggest that the socio-economical factors specified in the previous studies could also be influential in WNV infection development in our country.

The impact of educational status in getting information on the disease and on the transmission routes has also been reported (34). Also, Rios et al. demonstrated that WNV and Saint Louis Encephalitis (SLE) virus cause a greater number of neuroinvasive diseases in people with a low level of educational status (33). In the study group, it was detected that as the educational level decreased, WNV IgG seroprevalence increased significantly ($p<0.001$). This difference was prominent, especially in illiterate or merely literate people or elementary school graduates.

The strong aspect of the study is that a sample comprising all populations over two years of age from the whole province chosen by random sampling was included in the study. The sample size is relatively big and it is considered to be reflecting the community. Although the results that were found to be indeterminate in the study were evaluated as negative, the fact that the positive antibodies were not confirmed by a method such as PRNT due to the lack of financial support may have caused the seroprevalence to be detected higher than it actually is, and this constitutes a limitation. Since there were no population-based seroprevalence studies on other flaviviruses in the region where the study was conducted, a healthy comparison could not be made in terms of cross-reactivity. In addition, detection of non-neutralizing antibodies to the nucleocapsid antigen and failure to detect neutralizing antibodies to envelope glycoproteins by ELISA may result in an overestimation of antibody seroprevalence.

Because there was no information in the health histories of participants related to a neuroinvasive disease, the lack of information on how many in

this group with these risk factors developed neuro-invasive disease or death constitutes another limitation. Another limitation of the study is that the population densities were calculated roughly by dividing the district population by the district area. If a detailed population density map could have been created, a healthier analysis of whether there was a seroprevalence difference between the district centers which are thought to have a denser population and rural areas that presumably have a less dense population could have been made.

In conclusion, in the community based present study, which was conducted in the whole Manisa district, WNV IgG seroprevalence was detected to be 3.8% with ELISA method. In the study, no impact of the presence of an underlying disease, living in a rural area or population density was detected on seropositivity whereas an increase in age, low income and living in regions with a low altitude increased seropositivity significantly.

Ethical Approval: Dokuz Eylül University Non-Invasive Research Ethics Committee approved the study on August 13, 2018 with the decision number of “2018/17-07”

Informed Consent: N.A.

Peer-review: Externally peer-reviewed

Author Contributions: Concept – A.G., Ö.A.Ö., İ.M.A.Ö., K.E., B.Ü.; Design – A.N.E., M.E., A.G., Ö.A.Ö.; Supervision – K.E., B.Ü.; Fundings – A.N.E., M.E.; Materials – A.G., Ö.A.Ö., İ.M.A.Ö., B.Ü.; Data Collection and/or Processing – A.G., A.N.E., M.E., Ö.A.Ö., B.Ü.; Analysis and/or

Interpretation – A.G., B.Ü., Ö.A.Ö.; Literature Review – B.Ü., Ö.A.Ö., K.E., İ.M.A.Ö., A.G.; Writer – A.G., Ö.A.Ö., A.N.E., M.E. B.Ü.; Critical Reviews – Ö.A.Ö., İ.M.A.Ö., K.E., B.Ü., A.N.E., M.E.

Conflict of Interest: The authors declare no conflict of interest.

Financial Disclosure: There are no financial conflicts of interest to disclose.

Acknowledgment: We would like to thank Dokuz Eylül University Department of Scientific Research Project fund for supporting this research.

REFERENCES

- Hayes EB, Sejvar JJ, Zaki SR, Lanciotti RS, Bode AV, Campbell GL. Virology, pathology, and clinical manifestations of West Nile virus disease. *Emerg Infect Dis.* 2005;11(8):1174-1179.
- Smithburn KC, Hughes TP, Burke AW, Paul JH. A neurotropic virus isolated from the blood of a native of Uganda. *Am J Trop Med Hyg.* 1940;S1-20(4):471-92. [\[CrossRef\]](#)
- Richman DD, Whitley RJ, Hayden FG, editors. *Clinical Virology*. 3rd ed. Washington DC: ASM Press; 2009. p.1288.
- West Nile virus [Internet]. Geneva: World Health Organization (WHO). (October 3, 2017; cited November 22, 2021). Available from: <https://www.who.int/news-room/fact-sheets/detail/west-nile-virus>.
- Sampathkumar P. West Nile virus: epidemiology, clinical presentation, diagnosis, and prevention. *Mayo Clin Proc.* 2003;78(9):1137-43. [\[CrossRef\]](#)
- Kalaycioglu H, Korukluoglu G, Ozkul A, Oncul O, Tosun S, Karabay O, et al. Emergence of West Nile virus infections in humans in Turkey, 2010 to 2011. *Euro Surveill.* 2012;17(21):20182.
- Republic of Turkey Ministry of National Defence, General Directorate of Mapping (2018; cited November 22, 2021). Available from: <https://www.harita.gov.tr/il-ve-ilce-yuzol-cumleri>
- Turkish Statistical Institute (TUIK, TURKSTAT), Data Portal for Statistics, Population and Demography (2014; cited November 22, 2021). Available from: <https://data.tuik.gov.tr/Kategori/Get-Kategori?p=Nufus-ve-Demografi-109>
- Brault AC, Langevin SA, Ramey WN, Fang Y, Beasley DW, Barker CM, et al. Reduced avian virulence and viremia of West Nile virus isolates from Mexico and Texas. *Am J Trop Med Hyg.* 2011;85(4):758-67. [\[CrossRef\]](#)
- Chowdhury P, Khan SA, Dutta P, Topno R, Mahanta J. Characterization of West Nile virus (WNV) isolates from Assam, India: insights into the circulating WNV in northeastern India. *Comp Immunol Microbiol Infect Dis.* 2014;37(1):39-47. [\[CrossRef\]](#)
- Frost MJ, Zhang J, Edmonds JH, Prow NA, Gu X, Davis R, et al. Characterization of virulent West Nile virus Kunjin strain, Australia, 2011. *Emerg Infect Dis.* 2012;18(5):792-800. [\[CrossRef\]](#)
- Papa A, Bakonyi T, Xanthopoulou K, Vázquez A, Tenorio A, Nowotny N. Genetic characterization of West Nile virus lineage 2, Greece, 2010. *Emerg Infect Dis.* 2011;17(5):920-2. [\[CrossRef\]](#)
- Gray TJ, Webb CE. A review of the epidemiological and clinical aspects of West Nile virus. *Int J Gen Med.* 2014;7:193-203. [\[CrossRef\]](#)

- 14 Meshkat Z, Chinikar S, Shakeri M, Manavifar L, Moradi M, Mirshahabi H, et al. Prevalence of West Nile virus in Mashhad, Iran: A population-based study. *Asian Pac J Trop Med*. 2015;8(3):203-5. [\[CrossRef\]](#)
- 15 Soliman A, Mohareb E, Salman D, Saad M, Salama S, Fayed C, et al. Studies on West Nile virus infection in Egypt. *J Infect Public Health*. 2010;3(2):54-9. [\[CrossRef\]](#)
- 16 Hadjichristodoulou C, Pournaras S, Mavrouli M, Marka A, Tserkezou P, Baka A, et al. West Nile virus seroprevalence in the Greek population in 2013: A nationwide cross-sectional survey. *PLoS One*. 2015;10(11):e0143803. [\[CrossRef\]](#)
- 17 Hızal K, Yenicesu I, Erdal B, Yeşilyurt E, Fidan I, Kalkancı A, et al. [Investigation of West Nile virus seroprevalence in healthy blood donors]. *Mikrobiyol Bul*. 2010;44(3):425-30. Turkish.
- 18 Ayturan S, Aydoğan S, Ergünay K, Özcebe OI, Us D. [Investigation of West Nile virus seroprevalence in Hacettepe University Hospital blood donors and confirmation of the positive results by plaque reduction neutralization test]. *Mikrobiyol Bul*. 2011;45(1):113-24. Turkish.
- 19 Ergunay K, Saygan MB, Aydoğan S, Menemenlioglu D, Turan HM, Ozkul A, et al. West Nile virus seroprevalence in blood donors from Central Anatolia, Turkey. *Vector Borne Zoonotic Dis*. 2010;10(8):771-5. [\[CrossRef\]](#)
- 20 Ertuğrul-Örücü N, Yenicesu İ. [National Guidelines for Preparation, Use and Quality Assurance of Blood and Blood Components] [Internet]. Ankara: Republic of Turkey Ministry of Health & Türk Kızılay (2016; cited November 22, 2021). Turkish. Available from: https://www.kanver.org/Upload/Dosya/ulusal_kan_rehberi.pdf
- 21 West Nile virus infection. Annual epidemiological report for 2017 [Internet]. Stockholm: European Centre for Disease Prevention and Control (ECDC). (June 2019; cited November 22, 2021). Available from: https://www.ecdc.europa.eu/sites/default/files/documents/AER_for_2017-West-Nile-virus-infection.pdf
- 22 Karakoc ZC, Tuzuner BM, Ergonul O, Pierro A, et al. West Nile virus infection in the Mesopotamia region, Syria border of Turkey. *Vector Borne Zoonotic Dis*. 2013;13(10):739-43.
- 23 Gazi H, Ozkutuk N, Ecemis O, Atasoylu G, et al. Seroprevalence of West Nile virus, Crimean-Congo hemorrhagic fever virus, Francisella tularensis and Borrelia burgdorferi in rural population of Manisa, western Turkey. *J Vector Borne Dis*. 2016;53(2):112-7.
- 24 Manisa Province 2017 Environmental Status Report [Internet]. Manisa: Republic of Turkey Ministry of Environment, Urbanization and Climate Change, Branch Office Responsible for EIA and Environmental Permits. (2018; cited November 22, 2021). Turkish. Available from: <https://webdosya.csb.gov.tr/db/ced/icerikler/son1-manisa-20181001151132.pdf>
- 25 Muslu H, Kurt O, Ozbilgin A. [Evaluation of mosquito species (Diptera: Culicidae) identified in Manisa province according to their breeding sites and seasonal differences]. *Turkiye Parazitoloj Derg*. 2011;35(2):100-4. Turkish. [\[CrossRef\]](#)
- 26 Sıkı M, Gediz Deltası (İzmir Kuş Cenneti) kuşları. *Ekoloji Çevre Dergisi*. 2002; 11(44): 11 - 16.
- 27 Eisen L, Bolling BG, Blair CD, Beaty BJ, Moore CG. Mosquito species richness, composition, and abundance along habitat-climate-elevation gradients in the northern Colorado Front Range. *J Med Entomol*. 2008;45(4):800-11. [\[CrossRef\]](#)
- 28 Liu H, Weng Q. Environmental factors and risk areas of West Nile Virus in southern California, 2007–2009. *Environ Model Assess*. 2012;17(4):441-52. [\[CrossRef\]](#)
- 29 Gibbs SE, Wimberly MC, Madden M, Masour J, Yabsley MJ, Stallknecht DE. Factors affecting the geographic distribution of West Nile virus in Georgia, USA: 2002-2004. *Vector Borne Zoonotic Dis*. 2006;6(1):73-82. [\[CrossRef\]](#)
- 30 Lockaby G, Noori N, Morse W, Zipperer W, Kalin L, Governo R, et al. Climatic, ecological, and socioeconomic factors associated with West Nile virus incidence in Atlanta, Georgia, U.S.A. *J Vector Ecol*. 2016;41(2):232-43. [\[CrossRef\]](#)
- 31 Trawinski PR, Mackay DS. Identification of environmental covariates of West Nile virus vector mosquito population abundance. *Vector Borne Zoonotic Dis*. 2010;10(5):515-26. [\[CrossRef\]](#)
- 32 Harrigan RJ, Thomassen HA, Buermann W, Cummings RF, Kahn ME, Smith TB. Economic conditions predict prevalence of West Nile virus. *PLoS One*. 2010;5(11):e15437. [\[CrossRef\]](#)
- 33 Rios J, Hacker CS, Hailey CA, Parsons RE. Demographic and spatial analysis of West Nile virus and St. Louis encephalitis in Houston, Texas. *J of Am Mosq Control Assoc*. 2006;22(2):254-63. [\[CrossRef\]](#)
- 34 Diaz-Quijano FA, Martínez-Vega RA, Rodríguez-Morales AJ, Rojas-Calero RA, Luna-González ML, Díaz-Quijano RG. Association between the level of education and knowledge, attitudes and practices regarding dengue in the Caribbean region of Colombia. *BMC Public Health*. 2018;18(1):143. [\[CrossRef\]](#)